Kuo, C.-C., 1330
Kwiecinski, G. G., 926
Lacey, E. A., 1135, 1198
Laird, A., 1437
Lam, C., 691
Larivière, S., 937
Lee, L.-L., 1330
Lee, T. E., Jr., 791
Lee-Thorp, J. A., 471
Lemons, P. R., 989
Lenney Williams, C., 177
Leslie, D. M., Jr., 1279, 1399
Levesque, L. P., 1279
Light, J. E., 753
Liu, H., 1410
Liu, J., 644
Logan, M., 897
Lombardi, L., 26
Luna, F., 272
Lyons, S. K., 385
Mabry, K. E., 20
Macdonald, D. W., 1341, 1356
Macías, S., 1428
Marcus, W., 471
Marker, L. L., 840
Markowitz, T., 702
Mateo, J. M., 1163
Matsubayashi, H., 234
McCain, C. M., 967
McCue, E., 513
McElhinny, T. L., 1019
McGuire, B., 513
McShea, W. J., 627
Millesi, E., 615
Miquelle, D. G., 288
Mora, E. C., 1428
Moreno, S., 26
Morgan, G. S., 729
Morzillo, A. T., 553
Moscarella, R. A., 1300
Mote, K., 989
Murakami, O., 1228
Nams, V. O., 1374
Nassar, J. M., 106
Neiburger, E. J., 190
Nentwig, W., 571
Nichols, J. V., 1044
Nicholson, M. C., 553
Nogueira, M. R., 225
Okarma, H., 243
O'Neill, M. B., 117
Orrock, J., 627
Pace, R. M., III, 691
Pagels, J., 627
Parsons, J. L., 1399
Passey, B. H., 456
Patterson, B. D., 190
Pattini, A., 541
Pedrosa, E., 1437
Peltier, T. C., 915
Peracchi, A. L., 225
Perchellet, C. C., 989
Phillips, A. V., 524
Pierce, D. J., 81
Pillay N., 505
Poly, P. D., 369
Porter C. A., 791
Punzo, F. 1112, 1437
Quigley, H. B., 288
Read, A. J., 948
Reed, K., 471
Reeder, S. A., 809
Rezende, E. L., 311
Rhoads, D. D., 691
Ricca, M. A., 659
Rickart, E. A., 1443
Robinson, T. F., 471
Rodgers, B. E., 117
Rogers, D. S., 129
Roth, J. K., 487
Rowland, M. M., 92
Ruczyński, I., 243
Rushton, S., 1356
Salsbury, C. M.,, 299
Sánchez-Cordero, V., 547
Sankar, K., 981
Sarre, S. D., 1059
Schleich, C. E., 185
Schmidly, D. J., 159
Schmidt, K., 243
Schwartz, C. C., 718
Servín, J., 547
Shaw, C. N., 123
Shenk, T. M., 1456
Shields, G. F., 718
Sims, R. A., 197
Sleep, D. J. H., 216
Smirnov, E. N., 288
Smith, J. M., 471
Smith, K. G., 691
Smith, W. P., 1044
Śnieżko, S., 243
Solomon, N. G., 1135, 1182
Sponheimer, M., 471
Stafford, B. J., 263
Steele, T. E., 418
Steiner, C., 822
Sternberg, L. da S. L., 106
Studier, E. H., 926
Sumbera, R., 278
Swenson, J. E., 536
Swiderski, D. L., 861
Szykman, M., 1006
Tabaranza, B. R., Jr., 1443
Tague, R. G., 1456
Tang-Martínez, Z., 1212
Thayer, T. C., 487
Theuerkauf, J., 243
Thorington, R. W., Jr., 263
Tinker, M. T., 55
Torres, M. R., 541
Trejo, A., 652
Tristiani, H., 1228
Trolle, M., 607
Turner, J. P., 665
van Aarde, R. J., 851
Van Den Bussche, R. A., 791, 809, 1244
van der Merwe, N. J., 471
Van Horn, R. C., 1019
van Tuinen, M., 403, 444
Vander Wall, S. B., 487, 1089
Vassallo, A. I., 185
Veilleux, J. P., 1068
Veilleux, S. L., 1068
Villafuerte, R., 26
Viranta, S., 1267
Vitullo, A. D., 877
von Helversen, O., 886
Wales, B. C., 92
Wang, D.-H., 1410
Wang, Z.-W., 1410
Waser, P. M., 1031
Watanabe, H., 1228
Weckerly, F. W., 659
Weston, J. L., 996
Whitaker, J. O., Jr., 1068
White, B. N., 123
Wickliffe, J. K., 117
Wieczorek, J. R., 1198
Wilson, P. J., 123
Winkelmann, J. R., 561
Winter, Y., 886
Wisdom, M. J., 92
Wiseman, P. A., 1237
Woodman, N., 832
Woods, B. C., 903
Worthy, A. J., 665
Würsig, B., 702
Yamaguchi, N., 1341, 1356
Zahler, P., 480
Zenuto, R. R., 877
Zervanos, S. M., 299
Zhang, Y., 644
Ziv, Y., 1

SUBJECTS

(New names in *italic* type)

Abundance:
  Glaucomys sabrinus griseifrons, 1044
  Oryctolagus cuniculus, 26
  paleomammology and, 354
  predicting patterns from process-based simulation model, 1
Accretion, in suckling phyllostomid bats, 926
Acinonyx jubatus, 840
Actenomys, 185
Activity patterns:
  Canis lupus, 243
  Octodon degus, 311
  Oryctolagus cuniculus, 26

Rattus argentiventer, 1228
Aepyceros melampus, 456, 471
Aerial surveys:
Enhdra lutris, 55
Africa:
Acinonyx jubatus, 840
Aepyceros melampus, 456, 471
Alcelaphus (2 spp.), 456
Alcelaphus buselaphus, 471
Antidorcas marsupialis, 471
Cephalophus (2 spp.), 471
Cephalophus (7 spp.), 456
Connochaetes (2 spp.), 471
Connochaetes taurinus, 456
Crocuta crocuta, 1006, 1019
Damaliscus (2 spp.), 456, 471
Dasymys incomtus, 505
East African Bovidae, 456
Gazella (2 spp.), 456
Heliophobius argenteocinereus, 278
Hippotragus (2 spp.), 456, 471
Kobus (3 spp.), 471
Kobus ellipsiprymnus, 456
Madoqua (2 spp.), 456
Mastomys (2 spp.), 851
Neotragus (2 spp.), 456
Oryx gazella, 456, 471
Ourebia ourebi, 456, 471
Panthera leo, 190
Raphicerus campestris, 471
Redunca (2 sp.), 471
Redunca fulvorufula, 456
Sigmocerus lichtensteinii, 471
Southern African Bovidae, 471
Sylvicapra grimmia, 456, 471
Syncerus caffer, 418, 456, 471
Taurotragus oryx, 418, 456, 471
Tragelaphus (4 spp.), 471
Tragelaphus (5 spp.), 456
Age estimation:
Crocuta crocuta, 1019
Age structure:
in fossil assemblages, 418
Glaucomys sabrinus griseifrons, 1044
Spermophilus citellus, 615
Urocyon cinereoargenteus, 996
Agile wallaby, 197
Aging, and learning in least shrew, 1112
Agouti paca, 185
Akodon:
longipilis, 652
xanthorhinus, 652
Alarm call:
Myocastor coypus, 254
Alaska:
Enhdra lutris, 55
Glaucomys sabrinus griseifrons, 1044
Ovibos moschatus, 915
Alcelaphus:
buselaphus, 471
buselaphus cokei, 456
buselaphus jacksoni, 456
Alces alces, 123, 718
Aleutian Islands:
Enhdra lutris, 55
Allometry:
of baculum in Mustelidae, 673
Mastomys (2 spp.), 851
Sciurus niger, 861
Allozymes:
of tropical harvest mice, 129
American fur seal, 65
American mink, 1341, 1356
Amur tiger, 288
Anoura:
cultrata, 886
geoffroyi, 886
Antelope:
dwarf, 456
roan, 456
sable, 456
Antidorcas marsupialis, 471
Antilocapra americana, 418, 1237
Arctocephalus:
australis, 65, 524
forsteri, 65
Argentina:
Akodon (2 spp.), 652
Ctenomys (2 spp.), 1198
Ctenomys sociabilis, 403
Ctenomys talarum, 272, 877
Eligmodontia morgani, 653
Myocastor coypus, 254
Oligoryzomys longicaudatus, 653
Tympanoctomys barrerae, 541
Arizona:
Cynomys gunnisoni, 1244
Dipodomys spectabilis, 1031
Arkansas:
Ursus americanus, 691
Artibeus jamaicensis, 926
Arvicola terrestris, 1356
Auditory bulla:
in subterranean and surface-dwelling caviomorph rodents, 185
Auroch, 418
Australia:
Macropus (5 spp.), 197
Phascolarctos cinereus, 897
Austria:
Spermophilus citellus, 615
Axis axis, 981
Baculum, in Mustelidae, 673
Bank vole, 117, 1292
Banner-tailed kangaroo rat, 1031
Barking:
Vulpes velox, 1417
Basal metabolic rate:
Mustela (2 spp.), 497
Bats:
Anoura (2 spp.), 886
Artibeus jamaicensis, 926
call alteration, 205
Carollia perspicillata, 886
Chaerephon johorensis, 205
Cheiromeles torquatus, 205
Chiroderma (2 spp.), 225
Choeronycteria mexicana, 886
clutter tolerance, 216
Eptesicus fuscus, 216
fig-feeding, 225
Glossophaga (2 spp.), 886
Glossophaga longirostris, 106
Hesperoptenus blanfordi, 205
Hylonycteris underwoodi, 886
Lasionycteris noctivagans, 216
Lasiurus cinereus, 216
Leptonycteris curasoae, 106, 886
Lichonycteris obscura, 886
Lonchophylla robusta, 886
Lophostoma (5 spp.), 791
Macroglossuss minimus, 561
Mops mops, 205
Myotis lucifugus, 216
nectar-feeding bat dependence on CAM plants, 106
nutrients in suckling, 926
operational tongue length in nectar-feeding, 886
Phyllostomus discolor, 926
Pipistrellus stenopterus, 205
Pipistrellus subflavus, 1068
Primonatalus prattae, 729
Pteronotus quadridens, 1428
Synconycteris australis, 561
Tonatia (7 spp.), 791
Bears:
Ursus americanus, 81, 123, 594, 691
Ursus arctos, 418, 536
Behavior. *See also specific behaviors; specific species*
inferring from fossil record, 418
Macropus agilis, 197
Myocastor coypus, 254
reunion behavior in fur seals, 524
Belding's ground squirrels, 1163
Beluga whale, 123
Biodiversity:
in Appalachian mountain forest, 627
Biogeography:
Limnomys bryophilus, 1443
Natalidae, 729
Ochotona princeps, 37
Sorex (15 spp.), 144
Bison, steppe, 418
Bison priscus, 418
Black bear, 81, 123, 594, 691
Black-footed ferret, 497
Black wildebeest, 418
Blarina:
brevicauda, 627, 1374
carolinensis, 20
Blue wildebeest, 456
Body composition:
Ovibos moschatus, 915

Body mass/weight:
Cervus elaphus nannodes, 659
Cynomys (5 spp.), 1254
distribution in fossil mammals, 431
Glaucomys sabrinus griseifrons, 1044
Marmota flaviventris, 903
Odocoileus hemionus, 659
Ursus americanus, 594
Body reserve:
Microtus brandtii, 1410
Body size:
patterns of species diversity and, 1
Phocoena phocoena, 948
Body temperature:
Marmota monax, 299
Bongo, 456
Borneo:
Tragulus javanicus, 234
Boselaphus tragocamelus, 981
Bos primigenius, 418
Bottlenose dolphin, 665
Brandt's vole, 1410
Brazil:
Chiroderma (2 spp.), 225
Leopardus pardalis, 607
Breeding:
Canis lupus, 243
Cynomys (2 spp.), 1244
evolution of induced ovulation in North American carnivores, 937
Heliophobius argenteocinereus, 278
Mastomys (2 spp.), 851
Rattus argentiventer, 1228
Brown bear, 536
Brown hyena, 418
Brushtail possums, 1059
Buffalo:
Cape, 418, 456
Giant, 418
Burchell's zebra, 418
Burrows:
Tympanoctomys barrerae, 541
Bushbuck, 456
Cache pilferage, 487
California:
Canis latrans, 177
Cervus elaphus nannodes, 659
Odocoileus hemionus, 659
Ursus americanus, 594
Callorhinus ursinus, 55
Camera-trapping:
Leopardus pardalis, 607
Campbell's hamster, 1141
CAM plants, nectar-feeding bat dependence on, 106
Canada:
Blarina brevicauda, 1374
Lepus americanus, 1100
Microtus pennsylvanicus, 1376
Peromyscus (2 spp.), 1374
Sorex (5 spp.), 1374
Vulpes velox, 1417
Canis:
latrans, 123, 177, 547, 989
lupus, 123, 243, 418
lupus dingo, 197
Canler's mountain reedbuck, 456
Cape buffalo, 418, 456
Capture probability:
Arctocephalus forsteri, 65
Capture-recapture analysis:
Leopardus pardalis, 607
Carnivores. *See also specific species*
evolution of induced ovulation in North American, 937
ranges of Miocene, 1267
Carollia perspicillata, 886
Catch-removal trapping, of voles, 1292
Cavia pamparum, 185
Cenozoic:
body mass distribution in fossil mammals, 431
Central America:
Reithrodontomys (10 spp.), 129
Cephalophus:
diet of, 456
monticola, 471
natalensis, 471
Cervus:
elaphus, 418, 1076
elaphus nannodes, 659
elaphus roosevelti, 89
unicolor, 981
Chaerephon johorensis, 205
Chamois, 418
Cheetah, 840
Cheiromeles torquatus, 205
Chemical communication:
in rodents, 1141
Chile:
Octodon degus, 311
China:
Microtus brandtii, 1410
Myospalax fontanierii, 644
Chinchilla lanigera, 185
Chinkara, 981
Chipmunk, yellow pine, 487
Chiroderma:
doriae, 225
villosum, 225
Chiroptera:
Artibeus jamaicensis, 926
Chaerephon johorensis, 205
Cheiromeles torquatus, 205
Chiroderma (2 spp.), 225
Hesperoptenus blanfordi, 205
Lasionycteris noctivagans, 579
Lasiurus (3 spp.), 579
Mops mops, 205
Phyllostomus discolor, 926
Pipistrellus stenopterus, 205
Pipistrellus subflavus, 1068
Primonatalus prattae, 729
Pteronotus quadridens, 1428
Yangochiroptera, 809
Chital, 981
Choeronycteria mexicana, 886
Chornobyl:
shrews from, 117
Clethrionomys:
gapperi, 20, 627, 956, 1374
glareolus, 117, 1292
rufocanus, 1292
Climate change, 354, 385, 403
Colombian small-eared shrew, 832
Colorado:
Marmota flaviventris, 903
Zapus (2 spp.), 1456
Columbia:
Cryptotis colombiana, 832
Communication, chemical:
in rodents, 1141
Community structure:
predicting patterns of species diversity, 1
range shifts of Pleistocene mammals, 385
Competition:
intraspecific in American mink, 1341
predicting patterns from process-based simulation model, 1
ungulatesin dry tropical forest, 981
Virginia opossum and raccoon, 1279
Conepatus:
chinga, 159
humboldtii, 159
leuconotus, 159
leuconotus leuconotus, 159
leuconotus texensis, 159
mesoleucus, 159
mesoleucus figginsi, 159
mesoleucus filipensis, 159
mesoleucus fremonti, 159
mesoleucus mearnsi, 159
mesoleucus mesoleucus, 159
mesoleucus nelsoni, 159
mesoleucus nicaraguae, 159
mesoleucus sonoriensis, 159
mesoleucus telmalestes, 159
mesoleucus venaticus, 159
semistriatus, 159
Conflict, parent-offspring:
Arctocephalus australis, 524
Ursus arctos, 536
Connochaetes:
gnou, 418, 471
taurinus, 456, 471
Conservation:
Conepatus (17 spp. and subspp.), 159
Gulo gulo, 92
Mustela vison, 1356
Odocoileus virginianus, 1300
paleomammology, 347
Panthera leo, 190

Vulpes velox, 1417
Corridors:
effects on population density and habitat use, 20
Cotton mouse, 20
Cotton rat, 20, 1399
Cover characteristics:
Odocoileus virginianus dakotensis, 1316
Coyote, 123, 177, 547, 989
Coypu, 254
Crocuta crocuta, 1006, 1019
Cryptomys:
damarensis, 1198
hottentotus, 1198
Cryptotis:
colombiana, 832
medellinia, 832
nigrescens, 832
parva, 1112
thomasi, 832
Ctenomys:
australis, 185
azarae, 185
haigi, 1198
magellanicus, 185
occultus, 185
opimus, 185
perrensis, 185
porteousi, 185
pundti, 185
rionegrensis, 185
sociabilis, 185, 403, 1198
talarum, 185, 272, 877
tuconax, 185
tucumanus, 185
Cuba:
Pteronotus quadridens, 1428
Cynomys:
gunnisoni, 1244, 1254
leucurus, 1254
ludovicianus, 185, 1254
mexicanus, 1254
parvidens, 1244, 1254
Cytochrome *b* gene:
Marmosa murina, 822
Marmota (5 sp.), 369
Neomys fodiens, 117
Pappogeomys alcorni, 753
Sorex (6 spp.), 117
Sorex (15 spp.), 144
Damaliscus:
hunteri, 456
lunatus, 456, 471
pygarus, 471
Dasymys incomtus, 505
Deer:
Odocoileus hemionus, 659, 1076
Odocoileus virginianus, 123, 418, 1300
Tragulus javanicus, 234
Deer mouse, 20, 1364
Degu, 311
Delphinapterus leucas, 123
Demographics:
Glaucomys sabrinus griseifrons, 1044
Lagenorhynchus obscurus, 702
Mustela vison, 1341
Spermophilus citellus, 615
Urocyon cinereoargenteus, 996
Denning behavior:
Mustela vison, 1356
Trichosurus vulpecula, 1059
Ursus americanus, 594
Dental disease, 190
Desert habitat:
Dipodomys spectabilis, 1031
North American rodent species richness, 967
Tympanoctomys barrerae, 541
Destructive sampling, 1292
Didelphis virginiana, 1279, 1464
Diet:
East African Bovidae, 456
Eupetaurus cinereus, 480
nectar-feeding bat dependence on CAM plants, 106
Ovibos moschatus, 915
Peromyscus leucopus, 1381
Petaurista philippensis, 1330
Sigmodon hispidus, 1399
Southern African Bovidae, 471
Dikdik, 456
Dingo, 197
Dipodomys:
bullar volume in, 185
panamintinus, 487
spectabilis, 1031
Discriminant function analysis, 1456
Disease transmission:
Trichosurus vulpecula, 1059
Dispersal:
Alces alces, 718
Canis latrans, 177
Heliophobius argenteocinereus, 278
Panthera tigris altaica, 288
philopatry in rodents, 1182
species diversity and, 1
Spermophilus citellus, 615
Distribution:
Lasionycteris noctivagans, 579
Lasiurus (3 spp.), 579
Miocene carnivores, 1267
models, 594
Ursus americanus, 594
Diversity. *See also* Genetic variation; Species diversity
Marmosa murina, 822
taxonomic inflation, 431
Dolichotis patagonum, 185
Dolphin:
bottlenose, 665
dusky, 702
Duikers, 465
Dusky dolphin, 702
Dwarf antelope, 456
Eastern pipistrelles, 1068
Echolocation:
call alteration, 205
clutter tolerance and, 216
Malaysian aerial-feeding insectivorus bats, 205
variation of calls, 1428
Ecology:
in Appalachian mountain forest, 627
paleomammology, 347
social rodents, 1182, 1198
Tympanoctomys barrerae, 541
Ecomorphology:
Miocene carnivores, 1267
Eland, 418, 456, 471
Eligmodontia morgani, 653
Elk:
Rocky Mountain, 418, 1076
tule, 659
Energetics:
Marmota monax, 299
Energy demand:
Microtus brandtii, 1410
Peromyscus leucopus, 1381
England:
Mustela vison, 1341, 1356
Oryctolagus cuniculus, 1356
Enhdra lutris, 55
Eptesicus fuscus, 216
Equus burchelli, 418
Estivation:
Marmota monax, 299
Estrous cycle:
Dasymys incomtus, 505
Eubalaena:
australis, 123
glacialis, 123
Eumetopias jubatus, 55
Eupetaurus cinereus, 480
European ground squirrel, 615
European rabbit, 26
Evolution. *See also* Paleomammology
of baculum in Mustelidae, 673
in Carnivora, 1267
of induced ovulation in North American carnivores, 937
molecular clock, 444
paleomammology, 347
Extinction:
Miocene carnivores, 1267
Extirpation:
pika populations in the Great Basin, 37
wolverine populations, 92
Family breakup:
Ursus arctos, 536
Feeding behavior. *See also* Foraging
fig-feeding bats, 225
Felis concolor, 92
Ferret, black-footed, 497
Field vole, 1292
Fisher, 92
Florida:

Primonatalus prattae, 729
Flying squirrel:
Indian giant, 1330
Prince of Wales, 1044
woolly, 480
Food availability:
Canis latrans, 547
Octodon degus, 311
Petaurista philippensis, 1330
timing of conception in Phocoena phocoena, 948
Food supplementation:
Marmota flaviventris, 903
Ovibos moschatus, 915
Footprints, in tracking tube, 1374
Foraging:
East African Bovidae, 456
Macroglossuss minimus, 561
olfaction of granivorous rodents, 487
Petaurista leucogenys, 263
by rodents for buried seeds, olfaction and, 1089
Sigmodon hispidus, 1399
Tragulus javanicus, 234
Forest-cover type:
black bear home range and, 81
Fossil record:
body mass distribution in North American mammals, 431
combining genetic data with, 444
mortality profiles for inferring behavior from, 418
Fossils. *See also* Paleomammology
Primonatalus prattae, 729
Fox:
gray, 996
swift, 989, 1417
Fox squirrel, 861
Franciscana dolphin, 123
French Guiana:
Marmosa murina, 822
Functional morphology:
operational tongue length in nectar-feeding bats, 886
Fur seal, South American, 524
Gastrointestinal tract:
Peromyscus leucopus, 1381
Gazella:
bennetti, 981
granti, 456
thomsonii, 456
Gender:
molecular method for determination, 123
Gene flow:
Odocoileus virginianus, 1300
Genetic variation:
Alces alces, 718
Canis latrans, 177
Ctenomys sociabilis, 403
Lagenorhynchus obscurus, 702
Reihrodontomys (10 spp.), 129
Ursus americanus, 691
Geographic differentiation:
Marmota (5 sp.), 369
Geographic information system (GIS) use:
Crocuta crocuta, 1006
Geographic variation:
social rodents, 1212
Gerbil, 185
Gerenuk, 456
Giant buffalo, 418
Glaciation:
phylogeography of moose in North America, 718
range shifts of Pleistocene mammals, 385
Glaucomys sabrinus griseifrons, 1044
Gliding behavior:
of Japanese giant flying squirrels, 263
Global warming, 354
Glossophaga:
commissarisi, 886
longirostris, 106
soricina, 886
Glossophaginae, 886
Golden hamster, 1141
Golden mouse, 20, 553
Granivory, 487, 1089
Grant's gazelle, 456
Gray fox, 996
Gray seal, 65
Gray-sided vole, 1292
Gray wolf, 123
Great Basin:
black bear distribution, 594
extirpation of pika populations in, 37
Great Basin pocket mouse, 487
Greater kudu, 456
Grizzly bears, 418
Ground squirrel, 615, 1163
Group living, in rodents, 1198
Growth:
Acinoyx jubatus, 840
Marmota flaviventris, 903
Mastomys (2 spp.), 851
Ovibos moschatus, 915
suckling phyllostomid bats, 926
Gulo gulo, 92
Gunnison's prairie dog, 1244
Gut scaling, 659
Habitat fragmentation:
corridor effects on population density and habitat use, 20
niche shift by Virginia opossum and, 1279
Vulpes velox, 989
Habitat model:
Gulo gulo, 92
Habitat selection:
bat, 216
Clethrionomys gapperi, 956
Didelphis virginiana, 1279
Macropus agilis, 197
Odocoileus virginianus dakotensis, 1316
Pipistrellus subflavus, 1068
Habitat structure:
Oryctolagus cuniculus, 26
Habitat use:
bats, 216
Cervus elaphus, 1076
corridor effects, 20
Ctenomys (2 spp.), 1198
Dipodomys spectabilis, 1031
Mustela vison, 1356
Odocoileus hemionus, 1076
Rattus argentiventer, 1228
Tragulus javanicus, 234
ungulates in dry tropical forest, 981
Ursus americanus, 81
Vulpes velox, 989
Halichoerus grypus, 65
Hamster, 1141
Harbor porpoise, 948
Harbor seal, 123
Hare, snowshoe, 1100
Harvest mice, 129
Hearing:
in subterranean and surface-dwelling caviomorph rodents, 185
Heliophobius argenteocinereus, 278
Hesperoptenus blanfordi, 205
Heterocephalus glaber, 185, 1198
Hibernation:
Marmota flaviventris, 903
Marmota monax, 299
Hippotragus:
equinus, 456, 471
niger, 456, 471
Hirola, 456
Holocene:
Ctenomys sociabilis, 403
mammalian response climate change, 354
Home range:
Crocuta crocuta, 1006
Lepus americanus, 1100
Macroglossuss minimus, 561
Mustela vison, 1341, 1356
Ochrotomys nuttalli, 553
Rattus argentiventer, 1228
Tragulus javanicus, 234
Ursus americanus, 81
Vulpes velox, 989
House mouse, 1141
Human activity, effects on:
Canis lupus, 243
Ochotona princeps, 37
paleomammals, 418
Ursus americanus distribution, 594
Humpback right whale, 123
Hyaena brunnea, 418
Hydrochaeris hydrochaeris, 185
Hyena:

brown, 418
spotted, 1006, 1019
Hylonycteris underwoodi, 886
Idaho:
Gulo gulo, 92
Ideal-free distribution models, 594
Illinois:
Ochrotomys nuttalli, 553
Immigration:
Spermophilus citellus, 615
Impala, 456, 471
Inbreeding:
Canis latrans, 177
India:
Axis axis, 981
Boselaphus tragocamelus, 981
Cervus unicolor, 981
Gazella bennetti, 981
Indiana:
Pipistrellus subflavus, 1068
Indian giant flying squirrels, 1330
Individuality, of Vulpes velox vocalization, 1417
Indonesia:
Rattus argentiventer, 1228
Invasive species:
Mustela vison, 1356
Jackson's hartebeest, 456
Jaguar, 191
Japanese giant flying squirrels, 263
Jaw shape:
Sciurus niger, 861
Kangaroo rat, 185
Panamint, 487
Kentucky:
Peromyscus leucopus, 1381
Kenya:
Crocuta crocuta, 1006, 1019
Kerodon rupestris, 185
Kin recognition:
in rodents, 1141, 1163
Kin structure:
Lepus americanus, 1100
Kobus:
ellipsiprymnus, 456, 471
leche, 471
vardonii, 471
Kongoni, 456
Kudu:
greater, 456
lesser, 456
Lactation:
Microtus brandtii, 1410
Peromyscus leucopus, 1381
phyllostomid bats, 926
Lagenorhynchus obscurus, 702
Lagidium viscacia, 185
Lagomorpha:
Lepus americanus, 1100
Ochotona princeps, 37
Oryctolagus cuniculus, 26, 1356
Lagostomus maximus, 185
Landscape:
effects on small mammals, 20, 627
model for wolverines, 92
Lasionycteris noctivagans, 216, 579
Lasiurus:
blossevilli, 579
borealis, 579
cinereus, 216, 579
Learning, spatial:
Cryptotis parva, 1112
Least blossum bat, 561
Least shrew, 1112
Leopardus pardalis, 607
Leptokurtic dispersal, 718
Leptonycteris curasoae, 106, 886
Lepus americanus, 1100
Lesser kudu, 456
Lichonycteris obscura, 886
Life history:
Dasymys incomtus, 505
Heliophobius argenteocinereus, 278
social rodents, 1182
Limnomys:
bryophilus, 1443
sibuanus, 1443
Lions, 190
Litocranius walleri, 456
Litter size:
Panthera tigris altaica, 288
Locomotion:
Ctenomys talarum, 272
positional behavior of Japanese giant flying squirrels, 263
Lonchophylla robusta, 886
Longevity:
Miocene carnivores, 1267
Spermophilus citellus, 615
Long-haired mouse, 652
Lophostoma:
brasiliense, 791
carrikari, 791
evotis, 791
schulzi, 791
silvicolum, 791
Louisiana:
Ursus americanus, 691
Lynx canadensis, 92
Macroglossuss minimus, 561
Macropus:
agilis, 197
eugenii, 197
fuliginosus, 197
giganteus, 197
rufogriseus, 197
Madoqua:
guentheri, 456
kirkii, 456
Major histocompatibility complex (MHC), 1141
Malawi:
Heliophobius argenteocinereus, 278
Malaysia:
Chaerephon johorensis, 205
Cheiromeles torquatus, 205
Hesperoptenus blanfordi, 205
Mops mops, 205
Pipistrellus stenopterus, 205
Malnutrition:
effect on Petaurus breviceps learning, 1437
Management:
Canis latrans, 177
Ochotona princeps, 37
Trichosurus vulpecula, 1059
Mark-capture:
Arctocephalus forsteri, 65
Spermophilus citellus, 615
Ursus americanus, 81
Marmosa murina, 822
Marmota:
broweri, 369
caligata, 369
flaviventris, 369, 903
monax, 299, 369
vancouverensis, 369
Marsupials:
Didelphis virginiana, 1279, 1464
Macropus agilis, 197
Marmosa murina, 822
Petaurus breviceps, 1437
Phascolarctos cinereus, 897
Trichosurus vulpecula, 1059
Martes pennanti, 92
Mastomys:
coucha, 851
natalensis, 851
Mate choice, in rodents, 1141, 1163
Maternal behavior:
Arctocephalus australis, 524
Microtus ochrogaster, 513
Ursus arctos, 536
Mating:
Antilocapra americana, 1237
Ctenomys talarum, 877
evolution of induced ovulation in North American carnivores, 937
Myocastor coypus, 254
Phocoena phocoena, 948
social rodents, 1182
Meadow vole, 20, 1141, 1374
Mediterranean ecosystem:
Oryctolagus cuniculus and habitat structure, 26
Megaptera novaeangliae, 123
Meriones unguiculatus, 185
Merycism:
Phascolarctos cinereus, 897
Mesocricetus auratus, 1141
Mesopredator, 1279
Metapopulation theory, 627
Metatherian, 1464
Mexican prairie dog, 1254
Mexico:
Canis latrans, 547
Pappogeomys alcorni, 753
Pappogeomys bulleri alcorni, 753

Reithrodontomys (10 spp.), 129
Microcavia australis, 185
Microhabitat use:
Dipodomys spectabilis, 1031
niche shift by Virginia opossum, 1279
Ochrotomys nuttalli, 553
Microsatellite DNA studies:
Antilocapra americana, 1237
Canis latrans, 177
Cynomys (2 spp.), 1244
Lepus americanus, 1100
Ursus americanus, 691
Microsciurus flaviventer, 263
Microtus:
agrestis, 1292
arvalis, 571
brandtii, 1410
ochrogaster, 513, 1141
pennsylvanicus, 20, 1141, 1376
Migration:
Lasionycteris noctivagans, 579
Lasiurus (3 spp.), 579
Milk:
from phyllostomid bats, 926
Miller's long-tongued bat, 107
Mink, American, 1341, 1356
Miocene:
carnivores, ranges of, 1267
mammalian response to climate change, 354
Primonatalus prattae, 729
Mirounga leonina, 65
Mitochondrial DNA studies:
Alces alces, 718
Conepatus (3 spp.), 159
Lagenorhynchus obscurus, 702
Marmosa murina, 822
Odocoileus virginianus, 1300
Pappogeomys alcorni, 753
round-eared bats, 791
Sorex (15 spp.), 144
Yangochiroptera, 809
Molecular clock, 444
Molecular phylogenetics:
Yangochiroptera, 809
Molecular systematics:
Sorex (15 spp.), 144
Mole rats, 185, 278, 1198
Montana:
Antilocapra americana, 1237
Gulo gulo, 92
Montane alpine habitat:
Clethrionomys gapperi, 956
Ochotona princeps, 37
Moose, 123, 718
Mops mops, 205
Morphological variation:
of baculum in Mustelidae, 673
Marmosa murina, 822
Morphology:
Cryptotis colombiana, 832
Limnomys bryophilus, 1443
Southern African Bovidae, 471
Morphometrics:
Acinonyx jubatus, 840
Pappogeomys alcorni, 753
Sciurus niger, 861
Tursiops truncatus, 665
Zapus (2 spp.), 1456
Mortality profiles, inferring behavior from, 418
Mountain lion, 92
Mouse:
cotton, 20
deer, 20, 487, 1364
golden, 20, 553
Great Basin pocket, 487
harvest, 129
house, 1141
long-haired, 652
Mastomys (2 spp.), 851
old-field, 20
Preble's meadow jumping, 1456
silky desert, 652
western jumping, 1456
white-footed, 20, 1374, 1381
yellow-nosed, 652
Mouse-deer, 234
Movement:
daily and seasonal, of elk and mule deer, 1076
daily by coyotes, 547
migration of tree bats in North America, 579
Rattus argentiventer, 1228
running speed in least shrew, 1112
Mule deer, 659, 1076
Murine mouse opossum, 822
Muskoxen, 915
Mus musculus domesticus, 1141
Mustela:
eversmannii, 497
nigripes, 497
vison, 1341, 1356
Mustelidae, baculum of, 673
Myocastor coypus, 185, 254
Myospalax fontanierii, 644
Myotis lucifugus, 216
Namibia:
Acinonyx jubatus, 840
Nannospalax ehrenbergi, 185
Napaeozapus insignis, 627
Nectar-feeding bats, 886
Neomys fodiens, 117
Neotragus:
batesi, 456
moschatus, 456
Neotropics:
Chiroderma (2 spp.), 225
Nepotism, 1163
Nested-clade analysis, 702
Nesting habits:
Ochrotomys nuttalli, 553
Rattus argentiventer, 1228
social rodents, 1182
Neuroendocrinology, 1212
Nevada:
Dipodomys panamintinus, 487
Perognathus parvus, 487
Peromyscus maniculatus, 487
Tamias amoenus, 487
Ursus americanus, 594
New Zealand:
Lagenorhynchus obscurus, 702
Trichosurus vulpecula, 1059
New Zealand fur seal, 65
New Zealand sea lion, 65
Niche relationships:
shift by Virginia opossum after reduction in raccoons, 1279
ungulates in dry tropical forest, 981
Nilgai, 981
Nitrogen, dietary:
Sigmodon hispidus, 1399
Noctilionoidea, 809
North America. *See also* United States
Alces alces, 718
body mass distribution in fossil mammals, 431
Conepatus (17 spp. and subspp.), 159
Cynomys (5 spp.), 1254
desert rodents, 967
evolution of induced ovulation in carnivores from, 937
Lasionycteris noctivagans, 579
Lasiurus (3 spp.), 579
Lepus americanus, 1100
Marmota (5 sp.), 369
tree bats, 579
North Atlantic right whale, 123
Northern fur seal, 55, 65
Norway rat, 1141
Nuclear recombination activator gene-2 (*Rag2*):
round-eared bats, 791
Nutria, 254
Ocelot, 607
Ochotona princeps, 37
Ochrotomys nuttalli, 20, 553
Octodon degus, 311
Odocoileus:
hemionus, 659
virginianus, 123, 1300
virginianus dakotensis, 1316
Oklahoma:
Didelphis virginiana, 1279
Procyon lotor, 1279
Sigmodon hispidus, 1399
Olfaction:
in fur seal reunion behavior, 524
in kin recognition, 1163
rodent, 487, 1089, 1163
Oligoryzomys longicaudatus, 653
Ontogeny, postnatal:
Dasymys incomtus, 505
Opossum:
murine mouse, 822
Virginia, 1279, 1464
Orcinus orca, 55
Oregon:

Cervus elaphus, 1076
Gulo gulo, 92
Odocoileus hemionus, 1076
Organ size, 1381
Oribi, 456
Oryctolagus cuniculus, 26, 1356
Oryx gazella, 456, 471
Oryzomys palustris, 20
Os penis, in Mustelidae, 673
Osteology:
Cryptotis colombiana, 832
Ourebia ourebi, 456, 471
Ovibos moschatus, 915
Ovulation:
evolution of induced ovulation in North American carnivores, 937
Pakistan:
Eupetaurus cinereus, 480
Paleomammology:
behavior inferred from mortality profiles, 418
body mass distribution in fossil mammals, 431
extracting time from phylogenies, 444
global warming response, 354
interface with mammology, 347
marmots, 369
paleophylogeography, 369
range shifts of Pleistocene mammals, 385
tuco-tuco, 403
Panamint kangaroo rat, 487
Panthera:
leo, 190, 418
tigris altaica, 288
Pappogeomys:
alcorni, 753
bulleri alcorni, 753
Papua New Guinea:
Macroglossuss minimus, 561
Parental behavior:
Arctocephalus australis, 524
Microtus ochrogaster, 513
Ursus arctos, 536
Parturition:
Didelphis virginiana, 1464
Microtus ochrogaster, 513
Patagonian tuco-tuco, 1198
Paternal behavior:
Microtus ochrogaster, 513
Paternity, multiple:
Antilocarpa americana, 1244
Cynomys (2 spp.), 1244
PCR (polymerase chain reaction):
gender determination method, 123
Pelorovis antiquus, 418
Pelvis, sexual dimorphism in, 1464
Pennsylvania:
Marmota monax, 299
Perognathus parvus, 487
Peromyscus:
gossypinus, 20
leucopus, 20, 627, 1374, 1381
maniculatus, 20, 129, 487, 627, 1374
polionotus, 20
Peru:
Arctocephalus australis, 524
Pesticides, in soil, 1089
Petaurista:
leucogenys, 263
philippensis, 1330
Petaurus breviceps, 1437
Petrogale xanthopus, 197
Phascolarctos cinereus, 897
Pheromones, 1141
Philippines:
Limnomys (2 spp.), 1443
Philopatry, 1182
Phocarctos hookeri, 67
Phoca vitulina, 55
Phocoena phocoena, 948
Phodopus campbelli, 1141
Phyllostomidae, 886
Phyllostomus discolor, 926
Phylogeny:
Ctenomys (13 spp.), 185
extracting time from, 444
Marmota (5 sp.), 369
Mustelidae, 673
Natalidae, 729
shrews from Chornobyl, 117
Sorex (15 spp.), 144
tropical harvest mice, 129
Yangochiroptera, 809
Phylogeography:
Alces alces, 718
Odocoileus virginianus, 1300
paleophylogeography, 369
Pika, 37
Pinniped:
Arctocephalus (2 spp.), 65
Arctocephalus australis, 524
Callorhinus ursinus, 55, 65
Enhydra lutris, 55
Eumetopias jubatus, 55
Halichoerus grypus, 65
Mirounga leonina, 65
Phocarctos hookeri, 67
Phoca vitulina, 55, 123
Pipistrellus:
stenopterus, 205
subflavus, 1068
Plants. *See* Vegetation
Plateau zokors, 644
Pleistocene:
mammalian response to climate change, 354
range shifts, 385
Pocket gopher, 753
Poland:
Canis lupus, 243
Polecat, Siberian, 497
Pontoporia blainvillei, 123
Population bottleneck:
Alces alces, 718
Ctenomys sociabilis, 403
Population density:
Clethrionomys (2 sp.), 1292
corridor effects, 20
Enhydra lutris, 55
Leopardus pardalis, 607
Microtus agrestis, 1292
Spermophilus citellus, 615
tracking tube use for analysis, 1374
Population dynamics:
Dipodomys spectabilis, 1031
Lepus americanus, 1100
Spermophilus citellus, 615
Population genetics:
Ursus americanus, 691
Population structure:
Canis latrans, 177
Lagenorhynchus obscurus, 702
Odocoileus virginianus, 1300
Positional behavior:
Petaurista leucogenys, 263
Possum, brush tail, 1059
Postnatal development:
Heliophobius argenteocinereus, 278
Prairie dog:
black-tailed, 185, 1254
Gunnison's, 1244, 1254
Mexican, 1254
Utah, 1244, 1254
white-tailed, 1254
Prairie vole, 513, 1141
Preble's meadow jumping mouse, 1456
Predation:
by ancient hunters, 418
animal-human conflict, 190
avoidance by rodents, olfaction and, 1089
by Canis lupus, 243, 421
by Chiroderma (2 spp.) on figseeds, 225
of European rabbit in Mediterranean ecosystem, 26
of Macropus agilis, 197
mortality profiles for inferring behavior from fossil record, 418
by Mustela vison, 1356
by Phocoena phocoena, 948
of rodents by owls, 652
of Vulpes velox by coyote, 989
Primonatalus prattae, 729
Prince of Wales flying squirrel, 1044
Principal components analysis (PCA):
Conepatus (3 sp.), 159
Procyon lotor, 1279
Proechimys:
albispinus, 185
dimidiatus, 185
Pronghorn, 418, 1237
Protein malnutrition:

effect on Petaurus breviceps learning, 1437
Pteronotus quadridens, 1428
Quokkas, 197
Rabbit, 26, 1356
Raccoon, 1279
Radioactive fauna, 117
Radiotelemetry:
automatic tracking of small mammals, 571
Canis latrans, 547
Canis lupus, 243
Cervus elaphus, 1076
Crocuta crocuta, 1006
Lepus americanus, 1100
Macroglossuss minimus, 561
Marmota monax, 299
Mustela vison, 1341, 1356
Odocoileus hemionus, 1076
Panthera tigris altaica, 288
Phascolarctos cinereus, 897
Pipistrellus subflavus, 1068
Rattus argentiventer, 1228
Ursus arctos, 536
Vulpes velox, 989
Range. *See also* Home range
Miocene carnivores, 1267
North American desert rodents, 967
Odocoileus virginianus dakotensis, 1316
predicting patterns from process-based simulation model, 1
shifts of Pleistocene mammals, 385
Rangifer tarandus, 418
Raphicerus campestris, 471
Rat:
banner-tailed kangaroo, 1031
cotton, 20, 1399
Norway, 1141
red viscacha, 541
rice, 20, 652
ricefield, 1228
water, 505
Rattus:
argentiventer, 1228
norvegicus, 1141
Recombination activator gene-2 (*Rag2*):
round-eared bats, 791
Yangochiroptera, 809
Red-backed vole, 20, 627, 956, 1374
Red deer, 418
Red hartebeest, 471
Redunca:
arundinum, 471
fulvorufula, 456, 471
Red viscacha rat, 541
Reedbuck, Canler's mountain, 456
Reindeer, 418
Reithrodontomys:
chrysopsis, 129
creper, 129
fulvescens, 129
gracilis, 129
megalotis, 129
mexicanus, 129
microdon, 129
spectabilis, 129
sumichrasti, 129
tenuirostris, 129
Removal:
Canis latrans, 177
Reproduction:
Antilocapra americana, 1237
Canis lupus, 243
Crocuta crocuta, 1006
Cynomys (2 spp.), 1244
Dasymys incomtus, 505
evolution of induced ovulation in North American carnivores, 937
Glaucomys sabrinus griseifrons, 1044
Heliophobius argenteocinereus, 278
Microtus brandtii, 1410
Panthera tigris altaica, 288
Phocoena phocoena, 948
Urocyon cinereoargenteus, 996
Ursus americanus, 594
Resource partitioning:
bat, 216
Mustela vison, 1341
Odocoileus virginianus dakotensis, 1316
Reunion behavior:
Arctocephalus australis, 524
Ribosomal DNA studies:
Lophostoma (5 spp.), 791
Tonatia (7 spp.), 791
Ricefield rat, 1228
Rice rat, 20, 652
Road density:
Wolverine populations, effect on, 92
Roan antelope, 456
Rocky Mountain elk, 418, 1076
Rodentia. *See also* Social biology, of rodents
Agouti paca, 185
Akodon (2 spp.), 652
Blarina brevicauda, 627, 1374
Blarina carolinensis, 20
Cavia pamparum, 185
Clethrionomys gapperi, 20, 627, 956
Clethrionomys glareolus, 117
Cryptomys (2 spp.), 1198
Cryptotis (4 spp.), 832
Cryptotis parva, 1112
Ctenomys (13 spp.), 185
Ctenomys sociabilis, 403
Ctenomys talarum, 272, 877
Cynomys (2 spp.), 1244
Cynomys (5 spp.), 1254
Cynomys ludovicianus, 185
Dasymys incomtus, 505
Dipodomys panamintinus, 487
Dipodomys spectabilis, 1031
Dolichotis patagonum, 185
Eligmodontia morgani, 653
Eupetaurus cinereus, 480
Glaucomys sabrinus griseifrons, 1044
Heliophobius argenteocinereus, 278
Heterocephalus glaber, 185, 1198
Hydrochaeris hydrochaeris, 185
Kerodon rupestris, 185
Lagidium viscacia, 185
Lagostomus maximus, 185
Limnomys bryophilus, 1443
Marmota flaviventris, 903
Marmota monax, 299
Mastomys (2 spp.), 851
Mesocricetus auratus, 1141
Microcavia australis, 185
Microsciurus flaviventer, 263
Microtus brandtii, 1410
Microtus ochrogaster, 513, 1141
Microtus pennsylvanicus, 20, 1376
Mus musculus domesticus, 1141
Myocastor coypus, 185, 254
Myospalax fontanierii, 644
Nannospalax ehrenbergi, 185
Napaeozapus insignis, 627
Neomys fodiens, 117
North American desert rodents, 967
Ochrotomys nuttalli, 20, 553
Octodon degus, 311
olfaction by, 1089
Oligoryzomys longicaudatus, 653
Oryzomys palustris, 20
owl selection on size and sex classes of, 652
Pappogeomys alcorni, 753
Pappogeomys bulleri alcorni, 753
Perognathus parvus, 487
Peromyscus (2 spp.), 627, 1374
Peromyscus (4 spp.), 20
Peromyscus leucopus, 1381
Peromyscus maniculatus, 129, 487
Petaurista leucogenys, 263
Petaurista philippensis, 1330
Phodopus campbelli, 1141
Proechimys (2 spp.), 185
Rattus argentiventer, 1228
Rattus norvegicus, 1141
Reithrodontomys (10 spp.), 129
Sciurus igniventris, 263
Sciurus niger, 861
Sigmodon hispidus, 20, 1399
Sorex (3 spp.), 627
Sorex (5 spp.), 1374
Sorex (6 spp.), 117
Sorex (15 spp.), 117

Spermophilus citellus, 615
Tamias amoenus, 487
Tamias striatus, 627
Tympanoctomys barrerae, 541
Zapus (2 spp.), 1456
Roosevelt elk, 81
Roost sites:
Macroglossuss minimus, 561
Pipistrellus subflavus, 1068
Round-eared bats, 791
Rumen-reticulum capacity, 659
Rumination:
Phascolarctos cinereus, 897
Rupicapra rupicapra, 418
Russia:
Panthera tigris altaica, 288
Sable antelope, 456
Sambar, 981
Scent marking, 1141
Sciurus:
igniventris, 263
niger, 861
Seal:
American fur, 65
gray, 65
harbor, 123
New Zealand fur, 65
northern fur, 55, 65
South American fur, 524
southern elephant, 65
Sea lion:
New Zealand, 65
Steller, 55
Sea otters, 55
Seasonal distribution:
Lasionycteris noctivagans, 579
Lasiurus (3 spp.), 579
Seasonality:
Cervus elaphus, 1076
Cynomys (5 spp.), 1254
Heliophobius argenteocinereus, 278
Lagenorhynchus obscurus, 702
Mustela eversmannii, 497
of North American induced ovulation carnivores, 937
Odocoileus hemionus, 1076
Phocoena phocoena, 948
Seeds, olfaction of buried by rodents, 1089
Selection:
of rodents by owls, 652
Setonix brachyurus, 197
Sex ratio:
Glaucomys sabrinus griseifrons, 1044
Spermophilus citellus, 615
Urocyon cinereoargenteus, 996
Ursus americanus, 594
Sexual dimorphism:
Acinonyx jubatus, 840
Cynomys (5 spp.), 1254
Didelphis virginiana, 1464
Heliophobius argenteocinereus, 278
Mastomys (2 spp.), 851
Ovibos moschatus, 915
Tursiops truncatus, 665
Sexual segregation:
Odocoileus virginianus dakotensis, 1316
Sexual selection:
Cynomys (5 spp.), 1254
Short-tailed shrew, 1374
Shrew:
Blarina brevicauda, 1374
Blarina carolinensis, 20
Colombian small-eared, 832
Cryptotis parva, 1112
Neomys fodiens, 117
Sorex (5 spp.), 1374
Sorex (6 spp.), 117
Sorex (15 spp.), 144
Siberian polecat, 497
Siberian tiger, 288
Sigmocerus lichtensteinii, 471
Sigmodon hispidus, 20, 1399
Silky desert mouse, 652
Silvery mole rat, 278
Sitatunga, 456, 471
Skull morphology:
Limbomys (2 spp.), 1443
Tursiops truncatus, 665
Zapus (2 spp.), 1456
Skunk, white-backed hog-nosed, 159
Snap-trapping, of voles, 1292
Snowshoe hare, 1100
Social behavior:
Heliophobius argenteocinereus, 278
Lepus americanus, 1100
Social biology, of rodents:
chemical communication, 1141
ecology, 1198
kin recognition, 1163
philopatry, 1182
trends, challenges, and future directions, 1135, 1212
Social structure:
Crocuta crocuta, 1006
Myocastor coypus, 254
Rattus argentiventer, 1228
Soil:
pesticides in, 1089
plateau zokor effects on, 644
Sooty mustached bat, 1428
Sorex:
araneus, 117
arcticus, 1374
camtschatica, 144
cinereus, 144, 627, 1374
corona, 117
fumeus, 627, 1374
granarius, 117
haydeni, 144
hoyi, 144, 627, 1374
jacksoni, 144
longirostris, 144
minutus, 117
monticolus, 144
palustris, 144, 1374
portenkoi, 144
preblei, 144
pribilofensis, 144
samniticus, 117
tenellus, 144
trowbridgii, 144
tundrensi, 117
ugyunak, 144
vagrans, 144
South Africa:
Dasymys incomtus, 505
Mastomys (2 spp.), 851
South America:
Akodon (2 spp.), 652
Arctocephalus australis, 524
Chiroderma (2 spp.), 225
Cryptotis colombiana, 832
Ctenomys (2 spp.), 1198
Ctenomys sociabilis, 403
Ctenomys talarum, 272, 877
Eligmodontia morgani, 653
Glossophaga longirostris, 106
Leopardus pardalis, 607
Leptonycteris curasoae, 106
Marmosa murina, 822
Myocastor coypus, 254
Octodon degus, 311
Odocoileus virginianus, 1300
Oligoryzomys longicaudatus, 653
Tympanoctomys barrerae, 541
South American fur seal, 524
South Atlantic right whale, 123
South Carolina:
Peromyscus (2 spp.), 20
Sigmodon hispidus, 20
Urocyon cinereoargenteus, 996
South Dakota:
Odocoileus virginianus dakotensis, 1316
Southern elephant seal, 65
Southern long-nosed bat, 107
Space use:
Crocuta crocuta, 1006
Octodon degus, 311
tracking tubes for analysis of, 1374
Spacing behavior:
Lepus americanus, 1100
Mustela vison, 1341
Spatial learning:
Cryptotis parva, 1112
Species diversity:
North American desert rodents, 967
paleomammology and, 354
predicting patterns from process-based simulation model, 1
Species richness:
in mesic patches, 627
North American desert rodents, 967
Spermatozoa:

Antilocapra americana, 1237
Ctenomys talarum, 877
Spermophilus:
beldingi, 1163
citellus, 615
Spotted hyena, 1006, 1019
Squirrel:
Belding's ground, 1163
European ground, 615
fox, 861
Indian giant flying, 1330
Japanese giant flying, 263
Prince of Wales flying, 1044
tree, 263
woolly flying, 480
Stable isotope analysis:
East African Bovidae, 456
nectar-feeding bats, 106
Ovibos moschatus, 915
Southern African Bovidae, 471
Steenbok, 471
Stellar sea lion, 55
Steppe bison, 418
Stomach contents, analysis of:
Sigmodon hispidus, 1399
Subterranean rodents. *See also specific species*
Ctenomys (13 spp.), 185
Ctenomys talarum, 272, 877
sociality in, 1198
Sugar glider, 1437
Survival:
Arctocephalus forsteri, 65
Dipodomys spectabilis, 1031
Glaucomys sabrinus griseifrons, 1044
Marmota flaviventris, 903
Spermophilus citellus, 615
Urocyon cinereoargenteus, 996
Vulpes velox, 989
Sweden:
Clethrionomys (2 sp.), 1292
Microtus agrestis, 1292
Swift fox, 989, 1417
Sylvicapra grimmia, 456, 471
Syncerus caffer, 418, 456, 471
Synconycteris australis, 561
Systematics:
Pappogeomys alcorni, 753
round-eared bats, 791
Yangochiroptera, 809
Taiwan:
Petaurista philippensis, 1330
Tamias:
amoenus, 487
striatus, 627
Taurotragus oryx, 418, 456, 471
Taxonomy:
hog-nosed skunks, 159
inflation in North American fossil mammals, 431
Temperature, effects of:
global warming, 354
Octodon degus, 311
on olfaction of buried seeds by rodents, 1089
timing of conception in Phocoena phocoena, 948
Temporal variation:
Canis latrans, 177
Territoriality:
intrasexual in American mink, 1341
Macroglossuss minimus, 561
social rodents, 1182
Texas:
Vulpes velox, 989
Thomson's gazelle, 456
Tiger, 288
Timber harvest, effect on:
Clethrionomys gapperi, 956
Glaucomys sabrinus griseifrons, 1044
Time allocation:
Macropus agilis, 197
Tonatia:
bidens, 791
brasiliense, 791
carrikeri, 791
evotis, 791
saurophila, 791
schulzi, 791
silvicola, 791
Tongue length:
in nectar-feeding bats, 886
Tooth wear:
Crocuta crocuta, 1019
East African Bovidae, 456
in paleomammals, 418
Panthera leo, 190
Phascolarctos cinereus, 897
Sciurus niger, 861
Topi, 456
Torpor:
Marmota monax, 299
Tracking tube, 1374
Tragelaphus:
angasii, 471
eurycerus, 456
imberbis, 456
scriptus, 456, 471
spekii, 456, 471
strepsiceros, 456, 471
Tragulus javanicus, 234
Travel distances:
Canis latrans, 547
Tree bats, 579
Trichosurus vulpecula, 1059
Tuberculosis:
Trichosurus vulpecula, 1059
Tuco-tuco, 403, 877, 1198
Tule elk, 659
Tursiops truncatus, 665
Tympanoctomys barrerae, 541
Ukraine:
Neomys fodiens, 117
Sorex (2 spp.), 117
United Kingdom:
Mustela vison, 1341, 1356
Oryctolagus cuniculus, 1356
United States:
Antilocapra americana, 1237
Blarina brevicauda, 627
Canis latrans, 177
Cervus elaphus, 1076
Cervus elaphus nannodes, 659
Clethrionomys gapperi, 627, 956
Cynomys (2 spp.), 1244
Didelphis virginiana, 1279, 1464
Dipodomys panamintinus, 487
Dipodomys spectabilis, 1031
Enhydra lutris, 55
Felis concolor, 92
Glaucomys sabrinus griseifrons, 1044
Gulo gulo, 92
Marmota flaviventris, 903
Marmota monax, 299
Napaeozapus insignis, 627
Ochotona princeps, 37
Ochrotomys nuttalli, 553
Odocoileus hemionus, 659, 1076
Odocoileus virginianus dakotensis, 1316
Ovibos moschatus, 915
Perognathus parvus, 487
Peromyscus (2 spp.), 20, 627
Peromyscus leucopus, 1381
Peromyscus maniculatus, 487
Pipistrellus subflavus, 1068
Primonatalus prattae, 729
Procyon lotor, 1279
range shifts of Pleistocene mammals, 385
Sigmodon hispidus, 20, 1399
Sorex (3 spp.), 627
Tamias amoenus, 487
Tamias striatus, 627
Tursiops truncatus, 665
Urocyon cinereoargenteus, 996
Ursus americanus, 81, 594, 691
Vulpes velox, 989
Zapus (2 spp.), 1456
Urban interface:
Ursus americanus and, 594
Urocyon cinereoargenteus, 996
Ursus:
americanus, 81, 123
arctos, 418, 536
Utah:
Cynomys parvidens, 1244
Vegetation:
black bear home range and condition of, 81
cover characteristics for white-tailed deer, 1316
kangaroo rat population declines and, 1031
nectar-feeding bat dependence on CAM plants, 106
plateau zokor effects on, 644
roost sites of Pipistrellus subflavus, 1068

tongue length in nectar-feeding bats, 886
Venezuela:
Glossophaga longirostris, 106
Leptonycteris curasoae, 106
Odocoileus virginianus, 1300
Vespertilionoidea, 809
Virginia:
Blarina brevicauda, 627
Clethrionomys gapperi, 627
Napaeozapus insignis, 627
Peromyscus (2 spp.), 627
Sorex (3 spp.), 627
Tamias striatus, 627
Virginia opossum, 1279, 1464
Visual discrimination, in Petaurus breviceps, 1437
Vocalization:
in fur seal reunion behavior, 524
Vulpes velox, 1417
Vole:
bank, 117, 1292
Brandt's, 1410
common, 571
field, 1292
gray-sided, 1292
meadow, 20, 1141, 1374
prairie, 513, 1141
red-backed, 20, 627, 956, 1374
water, 1356
Vulpes velox, 989, 1417
Wallaby:
agile, 197197
Bennett's, 197
yellow-footed rock, 197
Washington:
Gulo gulo, 92
Ursus americanus, 81
Waterbuck, 456
Water rat, 505
Water vole, 1356
Weaning:
Ursus arctos, 536
Western jumping mouse, 1456
Whales:
Delphinapterus leucas, 123
Eubalaena (2 spp.), 123
Megaptera novaeangliae, 123
Pontoporia blainvillei, 123
White-backed hog-nosed skunk, 159
White-footed mouse, 20, 1374, 1381
White-tailed deer, 123, 423, 1300, 1316
White-tailed prairie dog, 1254
Wildebeest:
black, 418
blue, 456
Wolf, 123, 243, 418
Wolverine, 92
Woolly flying squirrel, 480
Wyoming:
Clethrionomys gapperi, 956
Odocoileus virginianus dakotensis, 1316
Zapus (2 spp.), 1456
Yellow-bellied marmot, 903
Yellow-nosed mouse, 652
Yellow pine chipmunk, 487
Zapus:
hudsonius preblei, 1456
princeps princeps, 1456
Zebra, Burchell's, 418
Zokor, plateau, 644